

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

**Prelim. Amdt dated November 21, 2003
REPLACEMENT SHEETS-DRAWINGS**

Figure 1

Delivery of Rho-antagonist tissue adhesive formulation.

1A. Application of tissue adhesive + Rho antagonist to the injured spinal cord with a pipette.



1B. Axon regeneration through the adhesion matrix + C3

Figure 2

Lesion of Corticospinal tract




Figure 3

**Corticospinal tract lesion
(untreated adult mice)**

Axon retraction

Lesion



BEST AVAILABLE COPY

Figure 4

Figure 4a




Figure 4b




Figure 4c





Figure 4d



BEST AVAILABLE COPY

Figure 5: Effect of C3/fibrin treatment on injured corticospinal tract

Figure 5a




Figure 5b



Figure 5c



BEST AVAILABLE COPY

Figure 6: C3/fibrin glue -treated spinal cord

Figure 6A




Figure 6B



BEST AVAILABLE COPY

Figure 7: Early Functional recovery

Figure 7A




Figure 7b



Figure 7C




Figure 8

BBB tests show recovery after C3 treatment

